

Flat Friedmann universe filled by dust and scalar field with multiple exponential potential

V R Gavrilov[§], V N Melnikov and S T Abdyrakhmanov

Centre for Gravitation and Fundamental Metrology, VNIIMS, and Institute for Gravitation and Cosmology, PFUR, 3-1 M.Ulyanovoy St., Moscow 117313, Russia

E-mail: gavr@rgs.phys.msu.su, melnikov@rgs.phys.msu.su and luxon@rgs.phys.msu.su

Abstract. We study a spatially flat Friedmann model containing a pressureless perfect fluid (dust) and a scalar field with an unbounded from below potential of the form $V(\varphi) = W_0 - V_0 \sinh\left(\sqrt{3/2}\kappa\varphi\right)$, where the parameters W_0 and V_0 are arbitrary and $\kappa = \sqrt{8\pi G_N} = M_p^{-1}$. The model is integrable and all exact solutions describe the recollapsing universe. The behavior of the model near both initial and final points of evolution is analyzed. The model is consistent with the observational parameters. We single out the exact solution with the present-day values of acceleration parameter $q_0 = 0.5$ and dark matter density parameter $\Omega_{\rho 0} = 0.3$ describing the evolution within the time approximately equal to $2H_0^{-1}$.

Submitted to: *Gen. Rel. Grav.*

PACS numbers: 04.20.Jb

[§] To whom correspondence should be addressed (gavr@rgs.phys.msu.su)

1. Introduction

Scalar fields play an essential role in modern cosmology. They are attributed to inflation models of the early universe and the models describing the present stage of the accelerated expansion as well. There is no unique candidate for the potential of the minimally coupled scalar field. Typically a potential is a sum of exponentials. Such potentials appear quite generically in a large class of theories (multidimensional [1] - [5], Kaluza-Klein models, supergravity and string/M - theories, see, for instance, [6], [7] and references therein) and from reconstruction and other schemes [8].

Single exponential potential was extensively studied within the Friedmann model using both qualitative methods [9] and exact solutions [10]. As to the multiple exponential potential, it's not studied well yet. Here we consider some multiple potential unbounded from below. According to [11] a Friedmann universe involving a scalar field with an unbounded from below potential exhibits the interesting features such as recollapsing even if it's flat.

In this paper we study the spatially flat Friedmann model containing a pressureless perfect fluid (dust) and a scalar field with an unbounded from below simplest multiple exponential potential of the form $V(\varphi) = W_0 - V_0 \sinh(\lambda\sqrt{8\pi G_N}\varphi)$, where the parameters W_0 and V_0 are arbitrary and $\lambda = \sqrt{3/2}$. For this value of λ the model is integrable and all exact solutions describe the recollapse of the universe within the finite time. The time as well as the intermediate behavior of the model crucially depend on parameters W_0 and V_0 . The paper is organized as follows. In section 2 we describe the model and obtain the equations of motion. The equations are integrated in section 3 using methods developed in our previous papers [1]-[3] and the general properties of these exact solutions are studied. In section 4 we fit the model to the observational parameters.

2. The model and equations of motion

We start with the spatially flat Friedmann universe filled by both a self-interacting scalar field φ with a potential $V(\varphi)$ and a separately conserved pressureless perfect fluid (dust). The corresponding Einstein equations read

$$H^2 = \frac{\kappa^2}{3} \left[\frac{1}{2} \dot{\varphi}^2 + V(\varphi) + \rho \right], \quad (1)$$

$$2\dot{H} + 3H^2 = -\kappa^2 \left[\frac{1}{2} \dot{\varphi}^2 - V(\varphi) \right], \quad (2)$$

where $H = \dot{a}/a$ is the Hubble parameter, ρ denotes the energy density of dust, overdot means derivative with respect to the cosmic time t , $\kappa = \sqrt{8\pi G_N} = M_p^{-1}$. Moreover, we have the equation of motion of the scalar field

$$\ddot{\varphi} = -3H\dot{\varphi} - V'(\varphi) \quad (3)$$

and the conservation equation for the perfect fluid

$$\dot{\rho} = -3H\rho. \quad (4)$$

The last equation gives immediately

$$\rho = \rho_0 \left(\frac{a_0}{a} \right)^3, \quad (5)$$

where the zero subindex means the present time, as usually. In what follows we consider the scalar field potential of the form

$$V(\varphi) = W_0 - V_0 \sinh \left(\sqrt{3/2} \kappa \varphi \right), \quad (6)$$

where W_0 and V_0 are arbitrary constants. As the system is symmetrical under the transformation $\varphi \rightarrow -\varphi$, $V_0 \rightarrow -V_0$, without a loss of generality we consider only the case $V_0 > 0$.

Now we introduce new variables x and y by the following transformation

$$a^3 = xy, \quad \kappa\varphi = \sqrt{2/3} \log(y/x), \quad x > 0, \quad y > 0. \quad (7)$$

Then the set of Eqs.(2),(3) results in

$$\ddot{x} = (\omega_1^2 - \omega_2^2)x - 2\omega_1\omega_2 y, \quad (8)$$

$$\ddot{y} = 2\omega_1\omega_2 x + (\omega_1^2 - \omega_2^2)y, \quad (9)$$

where we introduced the positive parameters ω_1 and ω_2 by

$$\omega_1^2 - \omega_2^2 = 3/4\kappa^2 W_0, \quad 2\omega_1\omega_2 = 3/4\kappa^2 V_0.$$

Then Eq.(1), where the presence of ρ is cancelled by Eq.(5), takes the form of the following constraint

$$\dot{x}\dot{y} - (\omega_1^2 - \omega_2^2)xy - \omega_1\omega_2(x^2 - y^2) = 3/4\kappa^2 \rho_0 a_0^3. \quad (10)$$

The set of Eqs.(8),(9) may be presented in the following complex form

$$\ddot{z} = \omega^2 z, \quad (11)$$

where we introduced the complex variable

$$z = x + iy$$

and the complex parameter

$$\omega = \omega_1 + i\omega_2.$$

It is easy to see that Eq.(11) implies the following complex integral of motion

$$\dot{z}^2 - \omega^2 z^2 = \text{const},$$

The presence of this complex integral is equivalent to the existence of 2 real integrals of motion. One of them $\Im(\dot{z}^2 - \omega^2 z^2) = -3/2\kappa^2 \rho_0 a_0^3$ represents the constraint given by Eq.(10) and the other integral $\Re(\dot{z}^2 - \omega^2 z^2)$ has an arbitrary value.

3. Exact solutions and behavior of the model

The equations of motion in form Eq.(11) are easily integrable. The result is

$$z = Ce^{i\alpha} \left(e^{\omega_1(t-t_0)+i\omega_2(t-t_0-\delta)} + e^{-\omega_1(t-t_0)-i\omega_2(t-t_0-\delta)} \right), \quad (12)$$

where $C > 0$. The constants t_0 and δ are arbitrary, C and α obey the relation

$$\Im \left(\omega^2 C^2 e^{i2\alpha} \right) = \frac{3}{8} \kappa^2 \rho_0 a_0^3$$

following from Eq.(10). Then substituting $x = \Re z$ and $y = \Im z$ into the relations given by Eq.(7) one easily gets the explicit expressions for the scale factor a and the scalar field φ .

Let us now consider the general properties of the exact solutions by analyzing the corresponding to Eq.(12) trajectories (orbits) of the moving point on the Cartesian xy plane. It follows from the definition of the variables x and y given by Eq.(7) that the physical segments of a trajectory belong to the angular domain with $x > 0$ and $y > 0$. Hereafter we show that each segments is of a finite length and its end-points are attached either to one coordinate axis or the both axes. From the physical viewpoint it means that all solutions describe the universe evolution within a finite time interval. Moreover, as the equations of motion are invariant with respect to the time reflection $t \rightarrow -t$ each segment of the trajectory may be passed in both directions. We notice that the constant factor $\exp(i\alpha)$ leads to the rotation of the trajectory about the origin $(0, 0)$ through the angle α . Further we consider only the case when $\alpha = 0$ taking into account that all remaining trajectories may be obtained by rotation.

If the time variable t is positive and large enough the last term in Eq.(12) is negligible ($\omega_1 > 0$). Then the motion is confined to the repelling logarithmic spiral with $x = C \exp[\omega_1(t-t_0)] \cos[\omega_2(t-T_0)]$ and $y = C \exp[\omega_1(t-t_0)] \sin[\omega_2(t-T_0)]$, where $T_0 \equiv t_0 + \delta$. As the time t grows the point clockwise rotates around the origin $(0, 0)$ and its distance from the origin increases as $\exp(\omega_1 t)$. It means that the evolution starts at some moment $t = T_0$ when the spiral intersects the axis x and finishes when it further intersects the axis y . The finite time of evolution is about the value $T = \pi/(2\omega_2)$. In this case the zeros of the functions $x(t)$ and $y(t)$ are different. Then near the initial point the expansion of the universe can be approximately described by the power-law equation $a(t) \sim (t - T_0)^{1/3}$. The similar equation $a(t) \sim [T - (t - T_0)]^{1/3}$ approximately describes the collapsing near the final point of evolution.

If $t < 0$ and $|t|$ is large enough the first term in Eq.(12) vanishes and the last term dominates. In this case the behavior of the model is similar with the only difference: the corresponding spiral is attracting. It describes the counterclockwise rotation with approaching to the origin of coordinates x and y .

The form of trajectories for the intermediate values of the time variable t is more complicated. To analyze the behavior one needs to superpose both attracting and repelling spirals. The result depends on the constant δ . Evidently it's enough to consider $\omega_2 \delta \in [0, \pi)$. If $\delta = 0$ we get $x = C \cosh[\omega_1(t-t_0)] \cos[\omega_2(t-t_0)]$ and $y = C \sinh[\omega_1(t-t_0)] \sin[\omega_2(t-t_0)]$. The corresponding trajectory with $C = 1$ is




Figure 1a




Figure 1b




Figure 1c




Figure 1d

Figure 1. The corresponding to the exact solutions trajectories on the xy plane. All remaining trajectories may be obtain by rotation about the origin $(0,0)$. Each solution is defined on some finite time interval which corresponds to a lying in the angular domain with $x > 0$ and $y > 0$ segment of the trajectory. End-points of such segment correspond to big bang and big crunch.

presented on Fig. 1a. The moving point passes the curve twice in both directions. It's not difficult to prove that for $\delta \neq 0$ any trajectory has no selfintersection points. On Figs. 1b and 1c the constant $\omega_2\delta$ ranges from 0 to $\pi/2$ and from $\pi/2$ to π , respectively. For $\omega_2\delta = \pi/2$ the trajectory passes through the origin center (see Fig. 1d).

This analysis shows that all exact solutions describe the evolution from big bang to big crunch within some finite time. Typically this time is about $T = \pi/(2\omega_2)$, however it may be arbitrarily shorter by choosing trajectories obtained after rotation. Each solution is defined on some finite time interval which corresponds to a lying in the angular domain with $x > 0$ and $y > 0$ segment of the trajectory. End-points of

such segment correspond to big bang and big crunch. If end-points of the segment are attached to different coordinate axes then the scalar field goes from $-\infty$ to $+\infty$ or vice versa during the evolution. If both end-points are attached to the coordinate axis y , the scalar field diverges to $+\infty$ as the time tends to the initial or final value. For almost all solutions the scale factor is proportional to the time in power $1/3$ near the initial and the final points of evolution. But if the end-point of the segment is in the origin $(0,0)$, the roots of the functions $x(t)$ and $y(t)$ coincide and the scale factor is proportional to the time in power $2/3$ near the corresponding point.

4. Fitting the model to observational parameters

After this qualitative description of the obtained exact solution (putting the present-day value of time t equal to zero) we present the solution to the equations of motion Eq.(11) in the form

$$z = (x_0 + iy_0) \cosh \omega t + \frac{\dot{x}_0 + i\dot{y}_0}{\omega} \sinh \omega t \quad (13)$$

with the values $x_0 \equiv x(0)$, $y_0 \equiv y(0)$, $\dot{x}_0 \equiv \dot{x}(0)$ and $\dot{y}_0 \equiv \dot{y}(0)$ expressed in the terms of the observational parameters: the Hubble constant H_0 , the present-day values q_0 and $\Omega_{\rho 0}$ of the acceleration parameter $q = a\ddot{a}/\dot{a}^2$ and the density parameter $\Omega_\rho = \kappa^2 \rho / (3H^2)$. One easily gets these expressions from Eqs.(1),(2) and the definition of the variables x and y given by Eq.(7)

$$x_0 = \left(\sqrt{1 + \varepsilon_0^2} + \varepsilon_0 \right)^{1/2} a_0^{3/2}, \quad (14)$$

$$y_0 = \left(\sqrt{1 + \varepsilon_0^2} - \varepsilon_0 \right)^{1/2} a_0^{3/2}, \quad (15)$$

$$\dot{x}_0 = \frac{3}{2} x_0 H_0 \left(1 \mp \sqrt{\frac{1}{3}(1 - q_0) - \frac{1}{2}\Omega_{\rho 0}} \right), \quad (16)$$

$$\dot{y}_0 = \frac{3}{2} y_0 H_0 \left(1 \pm \sqrt{\frac{1}{3}(1 - q_0) - \frac{1}{2}\Omega_{\rho 0}} \right), \quad (17)$$

where we denoted

$$\varepsilon_0 = \frac{9H_0^2}{8\omega_1\omega_2} \left[\frac{1}{3}(2 + q_0) - \frac{1}{2}\Omega_{\rho 0} \right] - \frac{\omega_1^2 - \omega_2^2}{2\omega_1\omega_2}. \quad (18)$$

The upper sign in Eqs.(16),(17) corresponds to the positive value $\dot{\varphi}_0 \equiv \dot{\varphi}(0)$, the lower sign appears when $\dot{\varphi}_0 < 0$. We notice that the model implies the value $(1 - q_0)/3 - \Omega_{\rho 0}/2$ to be nonnegative. It can be expressed via the present-day value $w_{\varphi 0}$ of the scalar field effective barotropic parameter

$$w_\varphi = \frac{p_\varphi}{\rho_\varphi} = \frac{\dot{\varphi}^2/2 - V(\varphi)}{\dot{\varphi}^2/2 + V(\varphi)}.$$

One easily obtain $(1 - q_0)/3 - \Omega_{\rho 0}/2 = (1 + w_{\varphi 0})(1 - \Omega_{\rho 0})/2 \geq 0$. Therefore the model is consistent with the observational data if the present-day value of the scalar field effective barotropic parameter is not less than -1 .




Figure 2a




Figure 2b




Figure 2c




Figure 2d

Figure 2. We present the exact solution for the following observational parameters: $q_0 = 0.5$, $\Omega_{\rho 0} = 0.3$ and the following model parameters: $\pi H_0/(2\omega_2) = 2$, $\varphi_0 = 0$, $\dot{\varphi}_0 > 0$.

The solution presented by Eqs.(13)-(17) with the scale factor a and the scalar field φ obtained from Eq.(7) exists in some finite time interval (t_{01}, t_{02}) with negative t_{01} and positive t_{02} where $x > 0$ and $y > 0$. Obviously, for given observational parameters H_0 , q_0 and $\Omega_{\rho 0}$ the values t_{01} and t_{02} and, consequently, the full time $(t_{02} - t_{01})$ of the universe evolution depends on parameters ω_1 and ω_2 determining the potential of the model. As we already mentioned in section 3, the typical time of evolution may be approximately estimated by the value $\pi/(2\omega_2)$. Then, instead of ω_1 and ω_2 we use further the following dimensionless model parameters: the typical time $\pi H_0/(2\omega_2)$ of the evolution in units of $H_0^{-1} \approx 14$ billions years and the present-day value $\kappa\varphi(0) \equiv \kappa\varphi_0 = \sqrt{2/3} \log(\sqrt{1 + \varepsilon_0^2} - \varepsilon_0)$ of the scalar field φ in units of $M_p = \kappa^{-1}$. Evidently, the model parameter ω_1 may be found for given ω_2 and φ_0 from Eq.(18). On Fig. 2 we present the exact solution for the following observational parameters: $q_0 = 0.5$, $\Omega_{\rho 0} = 0.3$, $\dot{\varphi}_0 > 0$ and the following model parameters: $\pi H_0/(2\omega_2) = 2$, $\varphi_0 = 0$. The interval of definition of the solution is turned




Figure 3. We introduce the following moments of time: t_{01} – big bang, t_{02} – big crunch, t_{03} – the beginning of the acceleration, t_{04} – the end of the accelerated stage of evolution, t_{05} – the beginning of the recollapsing.

to be with the following end-points $t_{01} \approx -0.6577H_0^{-1}$ and $t_{02} \approx 1.2222H_0^{-1}$. The scale factor presented on Fig. 2a may be described in the main order by $a \sim (t - t_{01})^{1/3}$ and $a \sim (t - t_{02})^{1/3}$ near the initial and final points. The scalar field presented on Fig. 2b diverges to $+\infty$ as $t \rightarrow t_{01}$ or $t \rightarrow t_{02}$. The scalar field potential given by Eq.(6) diverges to $-\infty$ as $t \rightarrow t_{01}$ or $t \rightarrow t_{02}$ (see Fig. 2c). Fig. 2d shows the domination of the scalar field near the initial and final singularities. Comparing Figs. 2c,2d one easily concludes that the kinetic term $\dot{\varphi}^2/2$ dominates the potential $V(\varphi)$ near these points. Then $w_\varphi \rightarrow 1$ as $t \rightarrow t_{01}$ or $t \rightarrow t_{02}$, i.e. the scalar field is like the stiff matter.

Now we analyze the behaviour of the model for various present-day values φ_0 of the scalar field. Besides, t_{01} and t_{02} , we introduce the following important moments of time:

- t_{03} – the beginning of the acceleration,
- t_{04} – the end of the accelerated stage of evolution,
- t_{05} – the beginning of the recollapsing (see Fig. 3).

Using the exact solution given by Eqs. (13)-(17) with the scale factor and the scalar field obtained from Eq. (7) for the parameter $q_0 = 0.5$, $\Omega_{\rho 0} = 0.3$ ($\dot{\varphi}_0 > 0$) and various φ_0 , we numerically calculate the mentioned time values. The result given in Table 1 shows the following property: the interval of the accelerated expansion as well as the full interval of evolution is shorter if the present-day value of the scalar field is greater. The corresponding behavior of the scalar factor is presented on Fig. 4.

$\varphi_0 \kappa$	$H_0 t_{01}$	$H_0 t_{03}$	$H_0 t_{04}$	$H_0 t_{05}$	$H_0 t_{02}$
-3	-0,6872	-0,2797	0,5645	2,1328	3,2765
-2,5	-0,6896	-0,2797	0,5645	2,1107	3,2054
-2	-0,6896	-0,2822	0,5718	2,0493	3,0213
-1,5	-0,6945	-0,2871	0,5865	1,9021	2,6728
-1	-0,7019	-0,2945	0,5988	1,6493	2,2162
-0,5	-0,7044	-0,2994	0,5301	1,3057	1,7229
0	-0,6577	-0,2675	0,3411	0,9056	1,2222
0,5	-0,5276	-0,1767	0,1546	0,5399	0,7780
1	-0,3607	-0,0834	0,0539	0,2847	0,4516
1,5	-0,2184	-0,0294	0,0171	0,1374	0,2503
2	-0,1251	-0,0073	0,0049	0,0638	0,1349
2,5	-0,0687	-0,0024	not found	0,0269	0,0711
3	-0,0368	not found	not found	0,0122	0,0368

Table 1. The dependence of the time values t_{01} , t_{02} , t_{03} , t_{04} and t_{05} explained on Fig. 3 on the present-day value φ_0 of the scalar field.




Figure 4. The scale factor for various present-day values φ_0 of the scalar field.

5. Conclusions

We studied the spatially flat Friedmann model containing a pressureless perfect fluid (dust) and a minimally coupled scalar field with an unbounded from below potential of the form $V(\varphi) = W_0 - V_0 \sinh\left(\lambda\sqrt{8\pi G_N}\varphi\right)$, where the parameters W_0 and V_0 are arbitrary. The parameter λ is chosen to be $\sqrt{3/2}$ in order to integrate the model in the explicit form. All exact solutions describe the recollapsing universe. The behavior of the model near both initial and final points of evolution is analyzed. Near the singularity the scale factor $a \sim t^{1/3}$ for almost all solutions. In this case the constraint Eq.(1) is dominated by the kinetic energy of the scalar field. Such solutions are called kinetic dominated [9]. There exists a special solution with $a \sim t^{2/3}$ near the singularity. It appears when the constraint Eq.(1) is dominated by the density of dust. It's so called fluid dominated solution [9]. The scalar field typically diverges to $\pm\infty$ as the time tends to the initial or final values. However, there is a special solution where the scalar field tends to some constant.

The evolution between big bang and big crunch crucially depends on the model parameters ω_1 and ω_2 . In particular, the time of evolution is determined by these parameters. We show that the model may be consistent with the observational parameters. We singled out the exact solution with the present-day values of the acceleration parameter $q_0 = 0.5$ and the dark matter density parameter $\Omega_{\rho 0} = 0.3$ describing the evolution within the time approximately equal to $2H_0^{-1}$.

Acknowledgments

This work was supported in part by the Russian Foundation for Basic Research (Grant 01-02-17312) and DFG Project 436 RUS 113/678/0-1. We are grateful to professor A.A. Starobinsky for helpful discussions. V.N.M. thanks Prof.Dr. Heinz Dehnen for the hospitality at the University of Konstanz during his stay there in October-November, 2003.

References

- [1] V. N. Melnikov, in: *Gravitation and Cosmology*, Proc. Int. Conf. on Gravitation and Cosmology (Ed. M. Novello), Rio de Janeiro 1993, (Singapore: Edition Frontieres 1994) p. 147.
- [2] V. N. Melnikov, *Multidimensional Classical and Quantum Cosmology and Gravitation: Exact Solutions and Variations of Constants*, CBPF-Notas de Fisica-051/93, Rio de Janeiro 1993, 93 pp.
- [3] V. N. Melnikov, in: *Gravitation and Cosmology II* (Ed. M. Novello), (Singapore: Edition Frontieres 1996), p. 465.
- [4] V. N. Melnikov, *Multidimensional Cosmology and Gravitation*, CBPF-MO-002/95, Rio de Janeiro 1995, 210 pp.
- [5] V. N. Melnikov, *Exact Solutions in Multidimensional Gravity and Cosmology III*, CBPF-MO-003/02, Rio de Janeiro 2002.
- [6] P. G. Ferreira and M. Joice, *Physical Review D* **58** 023503 (1998), astro-ph/9711102.
- [7] P. K. Townsend, *Quintessence from M-theory*, hep-th/0110072.
- [8] A. A. Starobinsky, *JETP Lett.* **68** 757 (1998).
- [9] I. P. C. Heard and D. Wands, *Cosmology with positive and negative exponential potentials*, gr-qc/0206085.
- [10] H. Dehnen, V. R. Gavrilo and Melnikov, *Grav. Cosm.* No. 3 (35) 189 (2003), gr-qc/0212107.
- [11] R. Kallosh and A. Linde, *Dark Energy and the Fate of the Universe*, astro-ph/0301087.